

The Rise and Fall of the Z-Pak: Updates in Antibiotic Guidelines for Common Urgent Care Conditions Worth Knowing

Joshua Russell, MD, MSc, ELS, FCUCM, FACEP

Senior Editor – EM:RAP, UCMax Podcast

Editor-in-Chief, *The Journal of Urgent Care Medicine (JUCM)*

Legacy-GoHealth Urgent Care

Metro Portland, Oregon

 CONTINUING EDUCATION COMPANY

1

The Rise and Fall of the Z-Pak:

Updates in Antibiotic Guidelines for Common Urgent Care Conditions Worth Knowing

Joshua Russell, MD, MSc, ELS, FCUCM, FACEP

Senior Editor – EM:RAP, UCMax Podcast

Editor-in-Chief, *The Journal of Urgent Care Medicine (JUCM)*

Legacy-GoHealth Urgent Care, Metro Portland, Oregon

2

Joshua Russell, MD
The Rise and Fall of the Z-Pak

Disclosure

I have no financial interests or relationships to disclose.



3

October 03, 2024



LETTER FROM THE EDITOR-IN-CHIEF

Antibiotic Prescribing in 'Gotham City'



I don't know about you, but I'm tired of being talked at about antibiotic stewardship. We all realize it's a problem. And though it's common practice to blame urgent care providers for the situation, we aren't uniquely culpable for antibiotic overuse.

To anyone paying attention, it's clear that antibiotic over-prescribing in urgent care remains as rampant as prima...

for the problems facing Gotham instead of the well-intentioned vigilante he is. The simplicity of having a single target of blame is tempting, but dangerous. Administrators, policy researchers and others who might speak on antibiotic stewardship without serving in the trenches risk invoking the same moral injury when they point the finger at UC clinicians as being solely responsible for antibiotic prescribing practices.

This is an important phenomenon to be considered as it...

Intermountain Health Receives Major Funding as Part of National Initiative to Advance Care for Children by Improving Antibiotic Prescribing Practices for Kids with Common Respiratory Tract Infections

Intermountain Health has been approved for a funding award by the Patient-Centered Outcomes Research Institute (PCORI) to help advance care for children by improving antibiotic prescribing practices for acute respiratory tract infection throughout its multi-state service area.

The \$2.5 million project will help Intermountain clinicians expand research-based practices for prescribing antibiotics to treat acute respiratory tract infection in children, avoid adverse side effects, and shrink the growing problem of drug-resistant infections.

4

Joshua Russell, MD
The Rise and Fall of the Z-Pak




Learning Objectives

1. Understand recent changes away from azithromycin in antibiotic guidelines for treatment of common bacterial infections encountered in Urgent Care
2. Develop familiarity with the few remaining indications of azithromycin
3. Identify the common and less common, but dangerous, adverse reactions associated with azithromycin

5

1. Antibiotic prescriptions per 1000 persons by state (sexiles) for all ages — United States,

Region	NUMBER OF ANTIBIOTIC PRESCRIPTIONS (MILLIONS)	ANTIBIOTIC PRESCRIPTIONS PER 1,000 PERSONS, RATE
Northeast	41.1	721
Midwest	49.5	719
South	106.2	825
West	39.5	502



Regional Outpt Antibiotic Prescribing

Source: CDC, 2022

6



Can we accurately gauge patient expectations?

7

INFECTIOUS DISEASE/ORIGINAL RESEARCH

Antibiotic Use for Emergency Department Patients With Upper Respiratory Infections: Prescribing Practices, Patient Expectations, and Patient Satisfaction

Samuel Ong, MD
Janet Nakase, MPH
Gregory J. Moran, MD
David J. Karras, MD
Matthew J. Kuehnert, MD
David A. Talan, MD
EMERGENCY ID NET Study Group

From Olive View—University of California Los Angeles Medical Center, Sylmar, CA (Ong, Nakase, Moran, Talan); Temple University School of Medicine, Philadelphia, PA (Karras); and the Centers for Disease Control and Prevention (Kuehnert).


Study objective: Physicians often prescribe antibiotics to patients even when there is no clear indication for their use. Previous studies examining antibiotic use in acute bronchitis and upper respiratory infections have been conducted in primary care settings. We evaluate the factors that physicians in the emergency department (ED) consider when prescribing antibiotics (eg, patient expectations) and the factors associated with patient satisfaction.

Methods: Ten academic EDs enrolled adults and children presenting with symptoms consistent with upper respiratory infection. Enrolled patients were interviewed before their physician encounter and were reinterviewed before discharge and 2 weeks later. Physicians were interviewed about factors that influenced their management decisions, including their perceptions of patients' expectations. Patients with a single diagnosis of uncomplicated acute bronchitis or upper respiratory infection were included for analysis.

Results: Of 272 patients enrolled, 68% of bronchitis patients and 9% of upper respiratory infection patients received antibiotics. Physicians were more likely to prescribe antibiotics when they believed that patients expected them (odds ratio [OR] 5.3; 95% confidence interval [CI] 2.9 to 9.6), although they were able to correctly identify only 27% of the patients who expected antibiotics. Satisfaction with the ED visit was reported by 87% of patients who received antibiotics and 89% of those not receiving antibiotics. Satisfaction with the visit was reported by 92% of patients who believed they had a better understanding of their illness but only by 72% of those who thought they had no better understanding (OR 4.4; 95% CI 2.0 to 8.4).

Conclusion: Physicians in our academic EDs prescribed antibiotics to 68% of acute bronchitis patients and to fewer than 10% of upper respiratory infection patients. Physicians were more likely to prescribe antibiotics to patients who they believed expected them, although they correctly identified only about 1 in 4 of those patients. Patient satisfaction was not related to receipt of antibiotics but was related to the belief they had a better understanding of their illness. [Ann Emerg Med. 2007;50:213-220.]

8



"Wallet with Cash" by 401(K) 2013 is licensed under CC BY-SA 2.0. To view a copy of this license, visit <https://creativecommons.org/licenses/by-sa/2.0/?ref=openverse>.

9

ORIGINAL INVESTIGATION

ONLINE FIRST

The Cost of Satisfaction

A National Study of Patient Satisfaction, Health Care Utilization, Expenditures, and Mortality

Joshua J. Fenton, MD, MPH; Anthony F. Jenant, MD; Klea D. Bertakis, MD, MPH; Peter Franks, MD

Background: Patient satisfaction is a widely used health care quality metric. However, the relationship between patient satisfaction and health care utilization, expenditures, and outcomes remains ill defined.

Methods: We conducted a **prospective cohort study of adult respondents (N=51,946)** to the 2000 through 2007 national Medical Expenditure Panel Survey, including 2 years of panel data for each patient and mortality follow-up data through December 31, 2006, for the 2000 through 2005 subsample (n=36,428). Year 1 patient satisfaction was assessed using 5 items from the Consumer Assessment of Health Plans Survey. We estimated the adjusted associations between year 1 patient satisfaction and year 2 health care utilization (any emergency department visits and any inpatient admissions), year 2 health care expenditures (total and for prescription drugs), and mortality during a mean follow-up duration of 3.9 years.

Results: Adjusting for sociodemographics, insurance status, availability of a usual source of care, chronic dis-

case burden, health status, and year 1 utilization and expenditures, **respondents in the highest patient satisfaction quartile** (relative to the lowest patient satisfaction quartile) had lower odds of any emergency department visit (adjusted odds ratio [aOR], 0.92; 95% CI, 0.84-1.00), higher odds of any inpatient admission (aOR, 1.12; 95% CI, 1.02-1.23), 8.8% (95% CI, 1.6%-16.6%) **greater total expenditures**, 9.1% (95% CI, 2.3%-16.4%) **greater prescription drug expenditures**, and **higher mortality** (adjusted hazard ratio, **1.05**; 95% CI, 1.03-1.33).

Conclusion: **In a nationally representative sample, higher patient satisfaction was associated with less emergency department use but with greater inpatient use, higher overall health care and prescription drug expenditures, and increased mortality.**

Arch Intern Med. 2012;172(5):405-411.
Published online February 13, 2012.
doi:10.1001/archinternmed.2011.1662



Scan for Audio
Audio Interview

W

HILE MOST HEALTH care quality metrics assess care processes and health outcomes, patient experience or satisfaction is considered a complementary measure of health care quality.¹ Patient satisfaction data may empower consumers to compare health plans and physicians,² and both the Centers for Medicare & Medicaid Services and the National Committee on Quality Assurance require participating health plans to publicly report patient satisfaction data.³ Health plans use patient satisfaction surveys to evaluate physicians and to determine incentive compensation, and consumer-oriented Web sites often report patient satisfaction ratings as the sole physician comparator.

Satisfied patients are more adherent to physician recommendations and more loyal to physicians,^{4,5} but research suggests a tenuous link between patient satisfaction and health care quality and outcomes.⁶⁻⁸ Among a vulnerable older population, patient satisfaction had no association with the technical quality of geriatric care,⁹ and evidence suggests that satisfaction has little or no correlation with Health Plan Employer Data and Information Set quality metrics.¹⁰

In addition, patients often request discretionary services that are of little or no medical benefit, and physicians frequently accede to these requests, which is associated with higher patient satisfaction.¹¹ Physicians whose compensation is more strongly linked with patient satisfaction are more likely to deliver discretionary services, such as advanced imaging for acute low back pain.¹²

See also page 435

See Invited Commentary at end of article

Author Affiliations: Department of Family and Community Medicine and Center for Healthcare Policy and Research, University of California-Davis, Sacramento.

10

Cognitive Costs of Abx Stewardship

JAMA Intern Med. 2014 December ; 174(12): 2029–2031. doi:10.1001/jamainternmed.2014.5225.

Time of Day and the Decision to Prescribe Antibiotics

Jeffrey A. Linder, MD, MPH, Jason N. Doctor, PhD, Mark W. Friedberg, MD, MPP, Harry Reyes Nieva, BA, Caroline Birks, MD, Daniella Meeker, PhD, and Craig R. Fox, PhD
Division of General Medicine and Primary Care, Brigham and Women's Hospital, Boston,

Results

There were 21 867 ARI visits to 204 clinicians in 23 practices that met our inclusion criteria; 44% resulted in antibiotic prescriptions (Table). Antibiotic prescribing increased throughout the morning and afternoon clinic sessions for antibiotics sometimes indicated and antibiotics never indicated ARIs (Figure). Relative to the first hour of a session, the adjusted odds ratios of antibiotic prescribing in the second, third, and fourth hours were 1.01 (95% CI, 0.91–1.13), 1.14 (95% CI, 1.02–1.27), and 1.26 (95% CI, 1.13–1.41), respectively ($P < .001$ for linear trend).

11



12

Automating the Common & Difficult Conversation


**SCRIPTING YOUR
LEAST FAVORITE
CONVERSATIONS**

**ANTIBIOTICS FOR
VIRAL URI**

Courtesy of Rob Orman, MD
(www.roborman.com/free-resources)

- Revise & Improve with Practice
- Save mental & emotional energy for MDM
- Start with top 2-3 most common painful conversations


13



Did COVID Usher in a
New Era of Antibiotic
Stewardship?

14

COVID-19 & Antibiotic Utilization



Tanislav C, Rosenbauer J, Kostev K. The COVID-19 Pandemic Enhanced the Decade-Long Trend of the Decreasing Utilization of Antibiotics. *Antibiotics*. 2023; 12(5):927.

15

PROVIDER SPECIALTY	NUMBER OF ANTIBIOTIC PRESCRIPTIONS (MILLIONS)	ANTIBIOTIC PRESCRIPTIONS PER PROVIDER, RATE
Primary Care Physicians	110.8	466
Physician Assistants and Nurse Practitioners	62.9	363
Surgical Specialties	19.5	219
Dentistry	25.1	205
Emergency Medicine	14.8	457
Dermatology	7.1	628
Obstetrics/Gynecology	6.3	167
Other	22.9	110
All Providers	269.4	295

Source: CDC, 2015


Table 3. Oral antibiotic prescribing by provider specialty — United States, 2020

Provider Specialty	Number of Antibiotic Prescriptions (Millions)	Antibiotic Prescriptions Per Provider, Rate
Primary Care Physicians	64.1	270
Physician Assistants & Nurse Practitioners	62.3	360
Surgical Specialties	15.3	172
Dentistry	23.4	191
Emergency Medicine	9.5	295
Dermatology	5.6	496
Obstetrics/Gynecology	4.6	123
Other	17.0	82
All Providers*	201.9	221

* Total may not add to all oral prescriptions (201.9 million) due to rounding.

16

Total Annual Azithromycin Prescriptions



17

Table 2. Top oral antibiotic classes and agents—United States, 2020

Characteristics: Antibiotic class	Number of Antibiotic Prescriptions (Millions)	Antibiotic Prescriptions Per 1,000 Persons, Rate
Penicillins	43.2	131
Cephalosporins	30.2	92
Macrolides	29	88
Tetracycline	22.7	69
B-lactams, increased activity	21	64
Characteristics: Antibiotic agent	Number of Antibiotic Prescriptions (Millions)	Antibiotic Prescriptions Per 1,000 Persons, Rate
Amoxicillin	39.3	119
Azithromycin	27.6	84
Amoxicillin/clavulanic acid	21	64
Cephalexin	19.6	60
Doxycycline	19.5	59

Source: CDC, 2020

ANTIBIOTIC CLASS	NUMBER OF ANTIBIOTIC PRESCRIPTIONS (MILLIONS)	ANTIBIOTIC PRESCRIPTIONS PER 1,000 PERSONS, RATE
Penicillins	53.2	159
Macrolides	36.1	108
Cephalosporins	36.0	108
B-lactams, increased activity	28.7	86
Tetracyclines	27.1	81
ANTIBIOTIC AGENT	NUMBER OF ANTIBIOTIC PRESCRIPTIONS (MILLIONS)	ANTIBIOTIC PRESCRIPTIONS PER 1,000 PERSONS, RATE
Amoxicillin	49.8	149
Azithromycin	34.9	105
Amoxicillin clavulanic acid	28.7	86
Doxycycline	24.1	72
Cephalexin	21.0	63

Source: CDC, 2022

18


Table 3. Oral antibiotic prescribing by provider specialty^a – United States, 2022

PROVIDER SPECIALTY	NUMBER OF ANTIBIOTIC PRESCRIPTIONS (MILLIONS)	ANTIBIOTIC PRESCRIPTIONS PER PROVIDER, RATE
Physician Assistants and Nurse Practitioners	84.4	165
Primary Care Physicians	70.0	179
Dentistry	25.2	127
Surgical Specialties	16.4	115
Emergency Medicine	12.0	182
Dermatology	5.4	299
Obstetrics/Gynecology	4.5	81
Other	18.5	27
All Healthcare Professionals	236.4	114

19

Sepsis Related Mortality By Year

Figure 1. Sepsis-related death rates for adults aged 65 and over, by age group: United States, 2000–2019



20

Pneumonia Related Mortality by Year

Year ↓	Deaths ↕	Population ↕	Crude Rate Per 100,000 ↕	Age Adjusted Rate Per 100,000 ↕
1999	63,730	279,040,168	22.8	23.5
2000	65,313	281,421,906	23.2	23.7
2001	62,034	284,968,955	21.8	22.2
2002	65,681	287,625,193	22.8	23.2
2003	65,163	290,107,933	22.5	22.6
2004	59,664	292,805,298	20.4	20.4
2005	63,001	295,516,599	21.3	21.0
2006	56,326	298,379,912	18.9	18.4
2007	52,717	301,231,207	17.5	16.8
2008	56,284	304,093,966	18.5	17.6
2009	53,692	306,771,529	17.5	16.5
2010	50,097	308,745,538	16.2	15.1
2011	53,826	311,591,917	17.3	15.7
2012	50,636	313,914,040	16.1	14.4
2013	56,979	316,128,839	18.0	15.9
2014	55,227	318,857,056	17.3	15.1
2015	57,062	321,418,820	17.8	15.2
2016	51,537	323,127,513	15.9	13.5
2017	55,672	325,719,178	17.1	14.3
2018	59,120	327,167,434	18.1	14.9
Total	1,153,761	6,088,633,001	18.9	17.7

21



22

The Rise of the Z-Pak...

- Synthetic Macrolide developed by Pliva Pharmaceuticals, Yugoslavia in 1981
- Licensed to Pfizer in 1991 – Z-Pak is born...
- Activity at 50S Ribosomal Subunit
- Clinically Favorable Profile
 - Delivered by phagocytes
 - Active at low pH
 - Long tissue half-life
- Rapidly became among top 5 most commonly prescribed antibiotics...

23

Azithromycin Resistance


- Population Level:
 - Rapid Emergence of Resistance of *S. pneumoniae* (& *S. pyogenes*) in 90's
 - Ribosomal Methylation (erm) or Efflux Pumping (mef)
 - Range of *S. PNA* Resistance from 10 – > 90% based on region
- Individual Level:
 - Asymptomatic Pneumococcal Carriage Rates up to 90%



24

Pneumococcal Resistance

Derek K-H. Ho et al, "Antibiotic Resistance in *Streptococcus pneumoniae* after Azithromycin Distribution for Trachoma", *Journal of Tropical Medicine*, vol. 2015, Article ID 917370, 8 pages, 2015.
<https://doi.org/10.1155/2015/917370>




25

”Gimme a Z-Pak to Knock It Down”

Common Z-Pak Requests:

- Sinusitis
- Cough/Bronchitis
- AOM - in PCN allergy
- Strep - in PCN allergy
- Chlamydia

26



HIGHLIGHTS OF PRESCRIBING INFORMATION
 These highlights do not include all the information needed to use ZITHROMAX® safely and effectively. See full prescribing information for ZITHROMAX.

ZITHROMAX (azithromycin) 250 mg and 500 mg tablets, for oral use
 ZITHROMAX (azithromycin) for oral suspension
 Initial U.S. Approval: 1991

----- RECENT MAJOR CHANGES -----

Warnings and Precautions, Cardiovascular Death (5.5) 11/2021

----- INDICATIONS AND USAGE -----

ZITHROMAX is a macrolide antibacterial drug indicated for mild to moderate infections caused by designated, susceptible bacteria:

- Acute bacterial exacerbations of chronic bronchitis in adults (1.1)
- Acute bacterial sinusitis in adults (1.1)
- Uncomplicated skin and skin structure infections in adults (1.1)
- Urethritis and cervicitis in adults (1.1)
- Genital ulcer disease in men (1.1)
- Acute otitis media in pediatric patients (6 months of age and older) (1.2)
- Community-acquired pneumonia in adults and pediatric patients (6 months of age and older) (1.1, 1.2)
- Pharyngitis/tonsillitis in adults and pediatric patients (2 years of age and older) (1.1, 1.2)

Limitation of Use:
 Azithromycin should not be used in patients with pneumonia who are judged to be inappropriate for oral therapy because of moderate to severe illness or risk factors. (1.3)
 To reduce the development of drug-resistant bacteria and maintain the effectiveness of ZITHROMAX (azithromycin) and other antibacterial drugs, ZITHROMAX (azithromycin) should be used only to treat infections that are proven or strongly suspected to be caused by susceptible bacteria. (1.4)

----- DOSAGE AND ADMINISTRATION -----

- **Adult Patients (2.1)**

Infection	Recommended Dose/Duration of Therapy
Community-acquired pneumonia (mild severity)	500 mg as a single dose on Day 1, followed by 250 mg once daily on Days 2 through 5.
Pharyngitis/tonsillitis (second-line therapy)	500 mg as a single dose on Day 1, followed by 250 mg once daily on Days 2 through 5.
Skin/skin structure (uncomplicated)	500 mg as a single dose on Day 1, followed by 250 mg once daily on Days 2 through 5 or 500 mg once daily for 3 days.
Acute bacterial exacerbations of chronic bronchitis (mild to moderate)	500 mg as a single dose on Day 1, followed by 250 mg once daily on Days 2 through 5 or 500 mg once daily for 3 days.
Acute bacterial sinusitis	500 mg once daily for 3 days.
Genital ulcer disease (chancroid)	One single 1 gram dose.
Non-gonococcal urethritis and cervicitis	One single 2 gram dose.
Gonococcal urethritis and cervicitis	One single 2 gram dose.

- **Pediatric Patients (2.2)**

Infection	Recommended Dose/Duration of Therapy
Acute otitis media (6 months of age and older)	30 mg/kg as a single dose or 10 mg/kg once daily for 3 days or 10 mg/kg as a single dose on Day 1 followed by 5 mg/kg/day on Days 2 through 5.
Acute bacterial sinusitis (6 months of age and older)	10 mg/kg once daily for 3 days.
Community-acquired pneumonia (6 months of age and older)	10 mg/kg as a single dose on Day 1 followed by 5 mg/kg once daily on Days 2 through 5.
Pharyngitis/tonsillitis (2 years of age and older)	12 mg/kg once daily for 5 days.

27

Acute Bacterial Sinusitis (ABRS)

Acute Sinusitis: Inflammation of nasal cavity & paranasal sinuses for <4 weeks

ABRS definition (per IDSA):

1. 10 days of illness **w/o improvement**
2. **Severe symptoms** (facial pain, purulent discharge) **AND** Fevers >39C x **3-4d**
3. **‘Double sickening’** (new onset fever, facial pain, headache after URI)

- <2% of cases are bacterial & **80%** of ABRS resolves w/I 2 weeks w/o ABX !!
- Higher Risk for Complicated Course: Advanced age, diabetes, immunosuppression

28

ABRS: What Do the Guidelines Say?

AAO-HNS (2015) | IDSA (2012) | ACP- CDC High Value Care Task Force (2016)

© 2022 UpToDate, Inc. and/or its affiliates. All Rights Reserved.

Distribution of pathogens in acute bacterial rhinosinusitis in adults

Pathogen	Incidence (%)
<i>Streptococcus pneumoniae</i>	20 to 43
<i>Haemophilus influenzae</i>	22 to 36
<i>Moraxella catarrhalis</i>	2 to 16
<i>Staphylococcus aureus</i>	10 to 13
<i>Streptococcus pyogenes</i>	3

Distribution of pathogens in acute bacterial rhinosinusitis based upon culture results.

Data from


1. Hooley JJ, Moayeri R, Decroix M, et al. Moxifloxacin five-day therapy versus placebo in acute bacterial rhinosinusitis. *Laryngoscope* 2010; 120:1027.
2. Rosenfeld RM, Piccirilli JJ, ChandraSekhar S, et al. Clinical practice guideline update: Adult sinusitis. *Otolaryngol Head Neck Surg* 2015; 152:51.

1. Treat only ABRS
2. Treat Immediately (IDSA) or Watch & Wait x 7 days (AAO-HNS)
 - Watchful waiting only if immunocompetent and good follow-up
3. **“Azithromycin/macrolides NOT recommended for empiric therapy due to high rates of *S. pneumoniae* resistance”**

ABRS: What Do the Guidelines Say?

- **Amoxicillin/Clavulanate 875mg BID** (NOT amoxicillin alone – Resistance in *H. flu* and *M. catarrhalis*)
- **Doxycycline 100mg BID** (PCN allergy/alternate first line)
- **Cefpodoxime 200mg BID**
- ****Levofloxacin 500mg daily** (only if can't tolerate alternatives; FQ risks)

Duration: 5-7 days



Community-Acquired Pneumonia (CAP)

Clinical diagnosis with constellation of findings: fever, dyspnea, cough, sputum production, abnormal lung sounds, abnormal CXR findings

- *'Clinicians should not perform testing or initiate antibiotic therapy in patients with bronchitis unless pneumonia is suspected'* -**ACP/CDC High Value Care Task Force, 2016**
- **Remember to educate:** Up to 3-4 weeks of cough is COMMON with bronchitis
- **Azithromycin is NOT recommended for outpatients (or inpatients) with COVID-19**



31

CAP: What Do the Guidelines Say?

2019 ATS/IDSA Joint Guideline Updates on Treatment of CAP:

- **Amoxicillin 1g TID -OR- Doxycycline 100mg BID x 5 days** (healthy, <65yo w/o recent antibiotic use)
- *Azithromycin/Macrolide ONLY Recommended if Local S. Pneumoniae resistance <25%*
- **Dual Therapy for >65 and/or co-morbidities and/or recent antibiotic use:**
 - Amox/clav -OR- 3rd gen cephalosporin **PLUS** Doxycycline -OR- Macrolide
 - **** Monotherapy with respiratory fluoroquinolones (e.g. levofloxacin) – least preferable option**

32

Penicillins in Urgent Care

Strep Pharyngitis

- Preferred first line: Penicillin VK (or Amoxicillin)
- *Macrolide resistant S. pyogenes*


Acute Otitis Media

- Preferred first line: Amoxicillin (+/- Clavulanate)
- *Macrolide resistant S. pneumoniae*
- Use **WAIT-AND-SEE** more

33

“I’m Allergic to Penicillin”

Penicillin is the most commonly reported drug allergy.¹



of patients in the US report penicillin allergy.¹

9 out of 10 reporting penicillin allergy are not truly allergic.⁴



80% of patients with IgE-mediated penicillin allergy lose the sensitivity after 10 years.⁴

Au LYC, Siu AM, Yamamoto LG. Cost and Risk Analysis of Lifelong Penicillin Allergy. *Clin Pediatr (Phila)*. 2019;58(11-12):1309-1314. doi:10.1177/0009922819853014

34

Consequences of 'Penicillin Allergy'

- Higher Lifelong Healthcare Spending
- Higher Rates of Broad Spectrum & Quinolone Abx Exposure
- Higher Rates of *C. difficile*




Au LYC, Siu AM, Yamamoto LG. Cost and Risk Analysis of Lifelong Penicillin Allergy. *Clin Pediatr (Phila)*. 2019;58(11-12):1309-1314. doi:10.1177/0009922819853014

35

Guidelines: De-Label When Able

Penicillin Risk Assessment

1. Adverse Reaction/Intolerance (e.g. GI upset) → PCN and Amoxicillin Safe
2. Mild/Delayed Hypersensitivity (e.g. maculopapular rash) → PO Test Dose of PCN or Cephalosporin
 - **Cephalosporin/Penicillin Cross Reactivity (mostly) Myth**
 - 97+% w/ true PCN allergy tolerate cephalosporins
 - **Consider Allergist Referral for Formal Testing**
1. Immediate/**IgE** Mediated (e.g. hives, anaphylaxis) → 3rd or later generation cephalosporin preferable, but all likely safe (0.001% risk of anaphylaxis)
2. Severe Non-Allergic Reaction (e.g. TEN, SJS, DRESS) → Avoid ALL B-lactams



36

De-Labeling Guidelines

Chua, KY et al. "The Penicillin Allergy Delabeling Program: A Multicenter Whole-of-Hospital Health Services Intervention and Comparative Effectiveness Study." *Clinical Infectious Diseases : an official publication of the Infectious Diseases Society of America* vol. 73,3 (2021): 487-496.

Dermatological		Respiratory or Systemic		Unknown	
Skin manifestation	Recommendation & Resultant allergy type	Clinical manifestation	Recommendation & Resultant allergy type	Clinical manifestation	Recommendation & Resultant allergy type
Childhood exanthem (unspecified) Mild rash with no severe features	<input type="checkbox"/> Unlikely to be significant (low severity)	Laryngeal involvement ("throat tightness" or "hoarse voice")	<input type="checkbox"/> Immediate hypersensitivity (low severity)	Unknown reaction ≤ 10 years ago	<input type="checkbox"/> Unknown (low severity)
Immediate diffuse rash ("itchy immediate rash") ≤ 2 hours post-dose	<input type="checkbox"/> Immediate hypersensitivity (low severity)	Respiratory compromise ("shortness of breath")	<input type="checkbox"/> Immediate hypersensitivity (low severity)	Unknown reaction > 10 years ago or family history of penicillin allergy only	<input type="checkbox"/> Unlikely to be significant (non-severe)
Diffuse rash or localized rash/swelling with no other symptoms (non-immediate or unknown timing)	> 10 years ago or unknown <input type="checkbox"/> Delayed hypersensitivity (low severity) ≤ 10 years ago <input type="checkbox"/> Delayed hypersensitivity (low severity)	Fever ("high temperature") Not explained by infection	<input type="checkbox"/> Delayed hypersensitivity (low severity)	Renal	
Angioedema ("to, facial or tongue swelling")	<input type="checkbox"/> Immediate hypersensitivity (severe)	Anaphylaxis or unexplained collapse	<input type="checkbox"/> Immediate hypersensitivity (severe)	Severe renal injury, failure or AIN (≥50% reduction in eGFR from baseline or absolute serum creatinine increase of ≥2.5mg/dL, or transplantation, or dialysis)	<input type="checkbox"/> Potential immune mediated (severe)
Generalized swelling (outside of angioedema)	<input type="checkbox"/> Immediate hypersensitivity (severe)	Haematological		Mild renal impairment (Does not meet criteria in box above)	<input type="checkbox"/> Unlikely immune mediated (non-severe)
Urticaria ("wheals and hives")	<input type="checkbox"/> Immediate hypersensitivity (low severity)	Low platelets ≤ 150 x10 ⁹ /L or unknown	<input type="checkbox"/> Potential immune mediated (severe)	Severe liver injury, failure or DILI (≥3x upper limit of normal (ULN) for ALT or AST, or ≥2x ULN for ALP with ≥2x ULN for bilirubin, or ≥2x ULN for ALP or transaminase)	<input type="checkbox"/> Potential immune mediated (severe)
Mucosal ulceration ("mouth, eye or genital ulcers")	<input type="checkbox"/> Delayed hypersensitivity (severe)	Low neutrophils ≤ 1x10 ⁹ /L or unknown	<input type="checkbox"/> Potential immune mediated (severe)	Mild hepatic enzyme derangement (Does not meet criteria in box above)	<input type="checkbox"/> Unlikely immune mediated (non-severe)
Pustular, blistering or desquamating rash ("skin shedding")	<input type="checkbox"/> Delayed hypersensitivity (severe)	Eosinophilia (≥0.7 x 10 ⁹ /L or unknown)	<input type="checkbox"/> Delayed hypersensitivity (severe)	Gastrointestinal, Neurological or Infusion-related	
Appropriate for supervised direct oral rechallenge (or direct de-labelling)			<input type="checkbox"/> Low risk	Severe neurological manifestation ("seizures or psychosis")	<input type="checkbox"/> Unknown or unclear mechanism
Appropriate for supervised direct oral rechallenge			<input type="checkbox"/> Low risk	Anaphylactoid/Infusion reaction (e.g. red man syndrome)	<input type="checkbox"/> Unknown or unclear mechanism
May be appropriate for referral for specialized skin testing			<input type="checkbox"/> Moderate risk		
May be appropriate for referral for specialized skin testing			<input type="checkbox"/> High risk		

37

Penicillin Allergy Risk : 'PEN-FAST'

PEN	Penicillin allergy reported by patient	<input type="checkbox"/> If yes, proceed with assessment
F	Five years or less since reaction ^a	<input type="checkbox"/> 2 points
A	Anaphylaxis or angioedema	<input type="checkbox"/> 2 points
S	Severe cutaneous adverse reaction ^b	
T	Treatment required for reaction ^a	<input type="checkbox"/> 1 point
		<input type="checkbox"/> Total points
Interpretation		
0	Very low risk of positive penicillin allergy test <1% (<1 in 100 patients reporting penicillin allergy)	
1-2	Low risk of positive penicillin allergy test 5% (1 in 20 patients)	
3	Moderate risk of positive penicillin allergy test 20% (1 in 5 patients)	
4-5	High risk of positive penicillin allergy test 50% (1 in 2 patients)	

Trubiano JA, Vogrin S, Chua KYL, et al. Development and Validation of a Penicillin Allergy Clinical Decision Rule. *JAMA Intern Med.* 2020;180(5):745-752. doi:10.1001/jamainternmed.2020.0403

38

PEN-FAST in Practice

- 73% of patients had PEN-FAST 0-2
- No patient with PEN-FAST <3 had +skin test
- 13% had safe Direct Oral Challenge

Performance of Each PEN-FAST Score in Predicting Penicillin Allergy

Performance Measure	Cutoff PEN-FAST score		
	0 vs 1-5	0-1 vs 2-5	0-2 vs 3-5
Sensitivity, % (95% CI)	100 (39.8-100)	100 (39.8-100)	100 (39.8-100)
Specificity, % (95% CI)	18.1 (11.6-26.3)	69.8 (60.6-78.0)	75.9 (67.0-83.3)

Su C, Belmont A, Liao J, Kuster JK, Trubiano JA, Kwah JH. Evaluating the PEN-FAST Clinical Decision-making Tool to Enhance Penicillin Allergy Delabeling. *JAMA Intern Med.* 2023;183(8):883-885. doi:10.1001/jamainternmed.2023.1572

Penicillin skin testing, No. (%)	104 (86.7)
Oral challenge, No. (%)	118 (98.3)
Oral challenge following skin test	102 (85.0)
Direct oral challenge	16 (13.3)
PEN-FAST score	
0	21 (17.5)
1	60 (50.0)
2	7 (5.8)
3	30 (25.0)
4	0
5	2 (1.7)

39

PEN-FAST Limitations

- Not Prospectively Validated in Adults (Bias?)
- **Higher Risk Features:** Rapid Onset of Reaction, Short Time Since Last Reaction, Requiring Hospitalization
- Not Valid in Children (i.e. < 18 years)

Copaescu AM, Vogrin S, Shand G, Ben-Shoshan M, Trubiano JA. Validation of the PEN-FAST Score in a Pediatric Population. *JAMA Netw Open.* 2022;5(9):e2233703

40

PEN-FAST - My Practice

- Don't use in Pts <18 yrs
- Use Shared Decision Making
- Direct Oral Challenge in Clinic if PF = 0-1
- If 2 or greater, refer to allergist
- Cephalosporins Generally Safe

41

Wait-and-See for AOM

- RCT of 283 children 6m – 12y seen in ED
- Abx filled: Immediate 62 % vs WASP 13%
- No difference in subsequent fever, otalgia, or revisit

Wait-and-See Prescription for the Treatment of Acute Otitis Media A Randomized Controlled Trial

David M. Spiro, MD, MPH
 Khoon-Yen Tay, MD
 Donald H. Arnold, MD, MPH
 James D. Dziura, PhD
 Mark D. Baker, MD
 Eugene D. Shapiro, MD

Context Acute otitis media (AOM) is the most common diagnosis for which antibiotics are prescribed for children. Previous trials that have evaluated a "wait-and-see prescription" (WASP) for antibiotics, with which parents are asked not to fill the prescription unless the child either is not better or is worse in 48 hours, have excluded children with severe AOM. None of these trials were conducted in an emergency department.

Objectives To determine whether treatment of AOM using a WASP significantly reduces use of antibiotics compared with a "standard prescription" (SP) and to evaluate the effects of this intervention on clinical symptoms and adverse outcomes related to antibiotic use.

Design, Setting, and Patients A randomized controlled trial conducted between July 12, 2004, and July 11, 2005. Children with AOM aged 6 months to 12 years seen in an emergency department were randomly assigned to receive either a WASP or an SP. All patients received ibuprofen and otic analgesic drops for use at home. A research assistant, blinded to group assignment, conducted structured phone interviews 4 to 6, 11 to 14, and 30 to 40 days after enrollment to determine outcomes.

Main Outcome Measures Filling of the antibiotic prescription and clinical course. **Results** Overall, 283 patients were randomized either to the WASP group (n=138) or the SP group (n=145). Substantially more parents in the WASP group did not fill the antibiotic prescription (62% vs 13%; P<.001). There was no statistically significant difference between the groups in the frequency of subsequent fever, otalgia, or unscheduled visits for medical care. Within the WASP group, both fever (relative risk [RR], 2.95; 95% confidence interval [CI], 1.75 - 4.99; P<.001) and otalgia (RR, 1.62; 95% CI, 1.26 - 2.03; P<.001) were associated with filling the prescription.

Conclusion The WASP approach substantially reduced unnecessary use of antibiotics in children with AOM seen in an emergency department and may be an alternative to routine use of antimicrobials for treatment of such children.

Trial Registration clinicaltrials.gov Identifier: NCT00250900
 JAMA. 2006;296:1235-1241 www.jama.com


ACUTE OTITIS MEDIA (AOM) is the most common reason for which an antibiotic is prescribed to children.^{1,2} Treatment of AOM accounts for an estimated 15 million antibiotic prescriptions written per year in the United States.³ Untreated AOM has a high rate of spontaneous resolution, with similar rates of complications whether antibiotics are prescribed or withheld.⁴⁻⁷ Resistance to antibiotics is a major public health concern worldwide and is associated with the widespread use of antibiotics.⁸⁻¹⁰

Two randomized trials have evaluated a management approach for AOM in which use of antibiotics is optional.^{11,12} One of these trials specifically evaluated children with nonsevere AOM,¹¹ while the other excluded

42

Wait-and-See Criteria

1. Non-severe sxs*
2. >6 months age
3. Unilateral
4. Can f/u w/i 2-3 days



43

Chlamydia

Chlamydia trachomatis - most common bacterial sexually transmitted genital infections

2nd Most Common Reportable Disease & 20% Inc since 2015

Intracellular Reproduction

Presentation ranges from Asymptomatic - > Dysuria/Discharge -> PID

44

Chlamydia: What Do The Guidelines Say?

CDC STI Treatment Guidelines, 2021

- Doxycycline 100mg BID x 7 days (**1st line**)
- 20% Failure Rate w/ Azithromycin 1g PO
- Doxy more effective for rectal and pharyngeal chlamydia infection
- Azithromycin 1g PO x 1 still preferred in pregnancy or if expect non-adherence with doxy
 - Consider “test of cure”

45

Toxicity and Adverse Reactions

- **Black box** : “Rare QTc prolongation and ventricular arrhythmias, including torsades de pointes”
- Gastrointestinal – Immediate (vomiting) & Delayed (diarrhea)
- Drug-Drug Interactions (CYP3A4)
- Liver Injury (rare, usually reversible, but fatal in cases if not stopped)

46

JAMA Network | Open.

Original Investigation | Pediatrics

Association of Inappropriate Outpatient Pediatric Antibiotic Prescriptions With Adverse Drug Events and Health Care Expenditures

Anne M. Butler, PhD, Derek S. Brown, PhD, Michael J. Durkin, MD, MPH, John M. Sahrmann, MA, Kaitlin B. Nickel, MPH, Caroline A. O'Neil, MA, MPH, Margaret A. Olsen, PhD, MPH, David Y. Hyun, MD, Rachel M. Zetts, MPH, Jason G. Newland, MD, MEd

Abstract

IMPORTANCE Inappropriate antibiotic prescribing for the treatment of pediatric infections is common, but the consequences of inappropriate antibiotics are not well described.

OBJECTIVE To evaluate the comparative safety and health care expenditures of inappropriate vs appropriate oral antibiotic prescriptions for common outpatient pediatric infections.

DESIGN, SETTING, AND PARTICIPANTS This cohort study included children aged 6 months to 17 years diagnosed with a bacterial infection (suppurative otitis media [SOM], pharyngitis, sinusitis) or viral infection (influenza, viral upper respiratory infection [URTI]), bronchiolitis, bronchitis, non-suppurative Otitis Media (NSOM) as an outpatient from April 1, 2016, to September 30, 2020, in the IBM MarketScan Commercial Database. Data were analyzed from August to November 2022.

EXPOSURES Inappropriate (ie, non-guideline-recommended) vs appropriate (ie, guideline-recommended) oral antibiotic agents dispensed from an outpatient pharmacy on the date of infection.

MAIN RESULTS AND MEASURES Proportionally score-weighted Cox proportional hazards models were used to estimate hazard ratios (HRs) and 95% CIs for the association between inappropriate antibiotic prescriptions and adverse drug events. Two-part models were used to calculate 30-day all-cause attributable health care expenditures by infection type. National level annual attributable expenditures were calculated by scaling attributable expenditures in the study cohort to the national outpatient population.

RESULTS The cohort included 2,042,240 eligible children (52% male, median [IQR] age, 6.14 [2.02] years). Overall, 37% to 38% received inappropriate antibiotics for bacterial infections and 6% to 10% for viral infections. Inappropriate antibiotics were associated with increased risk of several adverse drug events, including Clostridiaceae infection and severe allergic reaction among children treated with a nonrecommended antibiotic agent for a bacterial infection (among patients with suppurative Otitis Media [SOM], pharyngitis: HR, 6.23, 95% CI, 2.24-17.32; allergic reaction: HR, 4.14, 95% CI, 2.48-6.92). Thirty-day attributable health care expenditures were generally higher among children who received inappropriate antibiotics, ranging from \$21 to \$56 for bacterial infections and from ~\$96 to \$97 for viral infections. National-level attributable expenditure estimates were highest for suppurative Otitis Media (\$5.3 million), pharyngitis (\$2.1 million), and viral URIs (\$3.0 million).

CONCLUSIONS AND RELEVANCE In this cohort study of children with common infections treated in an outpatient setting, inappropriate antibiotic prescriptions were common and associated with increased risks of adverse drug events and higher attributable health care expenditures. These

(continued)

Open Access. This journal article distributed under the terms of the CC-BY License. JAMA Network Open. 2023;5(10):e232963. doi:10.1001/jamanetworkopen.2023.29263. May 16, 2023 | 1/6

JAMA Network | Open.

Original Investigation | Pediatrics

Association of Inappropriate Outpatient Pediatric Antibiotic Prescriptions With Adverse Drug Events and Health Care Expenditures


Anne M. Butler, PhD, Derek S. Brown, PhD, Michael J. Durkin, MD, MPH, John M. Sahrmann, MA, Kaitlin B. Nickel, MPH, Caroline A. O'Neil, MA, MPH, Margaret A. Olsen, PhD, MPH, David Y. Hyun, MD, Rachel M. Zetts, MPH, Jason G. Newland, MD, MEd

CONCLUSIONS AND RELEVANCE In this cohort study of children with common infections treated in an outpatient setting, inappropriate antibiotic prescriptions were common and associated with increased risks of adverse drug events and higher attributable health care expenditures. These

Inappropriate outpatient antibiotics increase health care utilization and spending in children

Medical Commentary
Supplemental content
Author affiliations and article information are listed at the end of this article.


47



When might Azithromycin be indicated?


1. Traveler's diarrhea
2. COPD Exacerbation
3. Atypical pneumonia
4. Chlamydia in Pregnancy or ?Adherence

48



NEW YORK
STATE OF OPPORTUNITY

Department of Health



NYC
Health

NEW YORK CITY DEPARTMENT OF HEALTH AND MENTAL HYGIENE

KATHY HOCHUL
Governor

JAMES V. McDONALD, M.D., M.P.H.
Commissioner

JOHANNE E. MORNE, M.S.
Executive Deputy Commissioner

DATE: 08/14/2024

TO: Healthcare Providers, Long Term Care Facilities, Healthcare Facilities, Clinical Laboratories, and Local Health Departments

FROM: New York State Department of Health (NYSDOH), New York City Department of Health and Mental Hygiene (NYC Health Department)

PROVIDER ADVISORY
Discontinue Use of Ciprofloxacin for Invasive Meningococcal Disease Post-Exposure Prophylaxis

Drug	Age	Dose	Duration	Other Considerations
Rifampin	<1 month	5 mg/kg, orally, every 12 hours	2 days	Consider consultation with a pediatric infectious disease expert for infants <1 month.
	≥1 month	10 mg/kg (maximum 600 mg), orally, every 12 hours	2 days	Can interfere with efficacy of oral contraceptives and some seizure prevention and anticoagulant medications; may stain soft contact lenses. Not recommended for pregnant people.
Ceftriaxone	<15 years	125 mg, intramuscularly	Single dose	To decrease pain at injection site, dilute with 1% lidocaine.
	≥15 years	250 mg, intramuscularly	Single dose	
Azithromycin	All Ages	10 mg/kg (maximum 500 mg)	Single dose	Alternative agent. In one study, equivalent to rifampin for eradication of <i>N. meningitidis</i> from nasopharynx.

Meningococcal Prophylaxis?

49

SUMMARY

- ❑ Patient satisfaction poorly correlated w/ Abx Rx
- ❑ Azithromycin stewardship improving...but still overprescribed in UC
- ❑ *S. PNA* & *S. Pyogenes* resistance is common

50



SUMMARY

- ❑ Azithro never recommended for ABRS
- ❑ Azithro monox is out for CAP
- ❑ PCN “allergy” can usually be de-labeled
- ❑ Wait-and-See for AOM
- ❑ Doxy is the way to go for Chlamydia
- ❑ Introduce the ‘Fall’ of Azithromycin to explain self-limited illness and risks