

Goodbye Pap, Hello Primary HPV Testing: Navigating the New Era of Cervical Cancer

David Glenn Weismiller, MD, ScM, FAAFP

Professor

Department of Family and Community Medicine
Kirk Kerkorian School of Medicine at the University of Nevada
Las Vegas, NV
david.weismiller@unlv.edu



1

Disclosure

I have no financial interests or relationships to disclose.




2

Learning Objectives

1. **Differentiate** between the three screening modalities: Primary HPV testing, Co-testing, and Cytology alone.
2. **Apply** the 2020 American Cancer Society (ACS) and updated USPSTF guidelines to determine the appropriate screening intervals for average-risk patients.
3. **Utilize** the ASCCP "Risk-Based Management" framework to determine when to "refer to colposcopy" versus "repeat in one year."
4. **Communicate** the rationale for less frequent screening to patients to alleviate anxiety regarding "missing" a cancer diagnosis.

Cervical Cancer



Surgical excision of cervix with a fungating squamous cell carcinoma

HPV and Cancer

Annual Estimates of HPV-Caused Cancers in the US

Among the more than 200 known HPV strains, **12–14 types are classified as "high-risk" (oncogenic)**. Two specific strains—**HPV 16 and HPV 18**—cause the vast majority of HPV-related malignancies across all anatomical sites.

Cancer Site	Total Cases in HPV-Associated Sites/Year	% caused by HPV	Dominant Strain in HPV (+) cancer (%)			Estimated Annual Cases Caused by HPV
			16	18	31, 33, 45, 52, 58	
Oropharynx (Head and Neck)	22, 585	~ 70%	90%	10%		16,000 (13, 600 men, 2,400 women)
Cervix	12, 287	~ 91%	55-60%	15-20%	20-30%	11,100
Anus	8, 348	~ 85-90%	85-90%	10-15%		7, 600 (5,300 women, 2,300 men)
Vulva	4, 370	~ 69%	55-60%	40-45%		3,000
Vagina	889	~ 75%	55-60%	40-45%		700
Penis	900					


(Note: While non-viral factors like tobacco and alcohol historically caused most head and neck cancers, **HPV 16 is now the primary driver of oropharyngeal cancers in North America**).

Impact of Vaccination: The CDC estimates that the 9-valent HPV vaccine can prevent roughly **92% (36,400)** of these annual HPV-caused cancers if administered prior to exposure.

National Center for Chronic Disease Prevention and Health Promotion;
Division of Cancer Prevention and Control

5



HPV 31 POSES A HIGHER RISK FOR CERVICAL PRECANCER AS COMPARED TO HPV 18



Stoler MH et al. Gynecol. Oncol. 2019; 153(1):26–33.

6

Cervical Cancer Progression Model



Schiffman, Mark H. et al. "Human papillomavirus and cervical cancer." *Lancet* 370 9590 (2007): 890-907.

7

The Clinical Conundrum 1

Patient: Sarah, a 31-year-old G2P2

History: No prior abnormalities; last screen 3 years ago was a "Normal" Pap (no HPV).

Today's Result: Primary HPV is **Positive** (Non-16/18). Reflex Cytology is **Negative** (NILM).

The Patient's Question: "Wait, you found the virus that causes cancer, but you say my cells are normal? Am I sick or not? And what do we do now?"

8

The Clinical Conundrum 2

Patient: Maria, a 31-year-old G2P2

History: No prior abnormalities; last screen 3 years ago was a "Normal" Pap (no HPV).

Today's Result: Primary HPV is **Positive for HPV16**.

The Patient's Question: *"Wait, you found the virus that causes cancer? Am I sick or not? And what do we do now?"*

9

The Clinical Conundrum 3

Patient: Sheila, a 57-year-old G3P3

- Establishing Care

- **History**

- LNMP – 4 years ago
- (+) VMS
 - Trial of Estrogen – no benefit
 - Recently had bioidentical Testosterone pellets implanted

- 8 years ago she underwent a hysterectomy for "precancer."
- She reports she has never had an abnormal pap and she has been getting "yearly evaluations for as long as she can remember."

The Patient's Question – *"Can I move to primary HPV testing instead of Pap?"*

10

Secondary Prevention

SCREENING FOR CERVICAL CANCER

11

The New Paradigm – Primary HPV Screening

- Cytology (the Pap smear) was a 20th-century triumph, but HPV testing is the 21st-century gold standard.
 - **The Change:** Major guidelines (ACS, ACOG, ASCCP) now prefer Primary HPV testing every 5 years starting at age 25.
 - **The Rationale:** HPV testing is significantly more sensitive than cytology. A negative HPV test provides greater reassurance against CIN 3+ over the next 10 years than a negative Pap alone.
 - **The "Exit" Strategy:** Screening can typically stop at age 65 if there is a "documented adequate negative prior screening" (3 consecutive negative paps or 2 negative HPV tests within 10 years).

12

Cervical Cancer (Dysplasia) Screening

Pap Smear

First reported 1928
(technique for
detecting uterine
cancer using
vaginal smears)



Dr. Pap (1883-1962)

Initially overlooked,
method gained traction
following collaborative
studies and publications -
1941

1950s: The test became widespread, drastically reducing
mortality rates.

13

Cervical Cancer (Dysplasia) Screening

Pap Smear

Pap Liquid Based Cytology

Introduced in 1996 following
FDA approval of the ThinPrep
System (SurePath-1999)

Standard in US by the 2010's

The method was introduced to improve upon conventional Pap
smears by decreasing the number of **inadequate samples**.

14

Cervical Cancer (Dysplasia) Screening

Pap Smear

Pap Liquid Based Cytology

(+) HPV Testing

- **2002:** The ACS included HPV testing as an adjunct (co-test) to cytology for women over 30.
- **2003:** The FDA approved the Hybrid Capture II test for co-testing.
- **2012:** Updated guidelines from organizations like ASCCP strengthened recommendations for co-testing every 5 years as a preferred method.

15

The New Paradigm Primary HPV Screening

2020 (ACS)

The Change: Major guidelines (ACS, ACOG, ASCCP) now prefer Primary HPV testing every 5 years starting at age 25.

The Rationale: HPV testing is significantly more sensitive than cytology. A negative HPV test provides greater reassurance against CIN 3+ over the next 10 years than a negative Pap alone.

The "Exit" Strategy: Screening can typically stop at age 65 if there is a "documented adequate negative prior screening" (3 consecutive negative paps or 2 negative HPV tests within 10 years).

**HPV testing is the 21st
Century Gold Standard**

16

Why the Shift?

From Cytology

The "Old" Logic (1950–2020): The Pap Smear/Cytology: We looked for "bad cells" (dysplasia) under a microscope.

- The Problem: It's a snapshot in time. It tells us if disease exists today, but it's not great at predicting who will get sick tomorrow.
- The **Sensitivity** Gap: A single Pap smear can miss up to 50% of high-grade lesions.

To Virology

The "New" Logic (2026 Paradigm):

- **The HPV Test:** Look for the **cause** (the virus) rather than the **symptom** (the cell change).
- **The Predictor:** HPV testing is significantly more **sensitive**.
 - (-) High-Risk HPV, risk of developing cervical cancer in the next 5–10 years is essentially zero.
- **The Efficiency:** Because the "Negative Predictive Value" is so high, we can safely extend screening intervals to every 5 years, reducing unnecessary biopsies and patient anxiety.

17

...a Long Way...

For 70 years, we played catch-up with the pap,

- trying to find abnormal cells after they already formed.

Today, we are playing defense. By testing for HPV,

- we identify the risk factor before the first abnormal cell ever appears.

We are not just finding precancer early anymore;

- we are preventing it from happening in the first place.

18

ARS Question #1


A 32-Year-Old Woman With No Prior History of Abnormal Cervical Cancer Screening Results Presents for a Wellness Examination. She Has Been Sexually Active With One Male Partner for the Past 10 Years and Has No Significant Medical History.

Which of the Following Statements Is True Regarding Screening for Cervical Cancer, Both in This Patient and in General?

- A. Following a hysterectomy (including the cervix) for endometrial cancer, women should undergo yearly vaginal cytology for 20 years.
- B. Current guidelines include cytology every 5 years as one screening option, beginning at age 21.
- C. Current guidelines recommend stopping screening at age 75
- D. The American Cancer Society recommends primary HPV testing every 5 years beginning at age 25.



19



- Cytology with Reflex HPV
- Cotesting
 - ☐ provides increased sensitivity and long-term negative predictive value of high-grade CIN and cervical cancer (CIN 3 or greater [CIN3+]) compared with cytology alone.
 - ☐ HPV testing is not recommended for individuals younger than 25 years because transient HPV infections are common in this group and rarely contribute to carcinogenic outcomes.
- Primary HPV Screening

Cervical Cancer Screening Tests

Fontham ETH, Wolf AMD, Church TR, et al. Cervical cancer screening for individuals at average risk: 2020 guideline update from the American Cancer Society. CA Cancer J Clin 2020;70:321–46.

20

Primary HPV Screening

- **Clinician-collected** primary HPV screening obtains a cervical cytology sample, which can first be tested for HPV.
- Depending on the testing method screening test reports positive based on the detection of high-risk type HPV messenger RNA E6/E7 proteins or DNA

Diagnostic Accuracy for Detection of Cervical Precancer (CIN2+) of Clinician-Collected Cervical Cytology, HPV Testing Based on Clinician-Collected Cervical Specimens (Clinician HPV), and HPV Testing Using Self-Collected Vaginal Specimens (Self-HPV)

	Pooled Sensitivity	Pooled Specificity
Cytology	80.4 (95% CI = 73.2-86.1)	78.5 (95% CI 69.8-85.2)
Clinician-HPV (PCR)	92.9 (95%CI = 88.6-95.5)	61.2 (95% CI = 41.2-78.1)

Fontham ETH, Wolf AMD, Church TR, et al. Cervical cancer screening for individuals at average risk: 2020 guideline update from the American Cancer Society. CA Cancer J Clin 2020;70:321-46.

21

Not statistically significant

Primary HPV Screening

- Clinician-collected primary HPV screening obtains a cervical cytology sample, which can first be tested for HPV.
- Depending on the testing method screening test reports positive based on the detection of high-risk type HPV messenger RNA E6/E7 proteins or DNA

Diagnostic Accuracy for Detection of Cervical Precancer (CIN2+) of Clinician-Collected Cervical Cytology, HPV Testing Based on Clinician-Collected Cervical Specimens (Clinician HPV), and HPV Testing Using Self-Collected Vaginal Specimens (Self-HPV)

	Pooled Sensitivity	Pooled Specificity
Cytology	80.4 (95% CI = 73.2-86.1)	78.5 (95% CI 69.8-85.2)
Self-HPV (PCR)	89.7 (95%CI 84.2-93.5)	64.7 (95% CI 44.6-80.7)
Clinician-HPV (PCR)	92.9 (95%CI = 88.6-95.5)	61.2 (95% CI = 41.2-78.1)

Fontham ETH, Wolf AMD, Church TR, et al. Cervical cancer screening for individuals at average risk: 2020 guideline update from the American Cancer Society. CA Cancer J Clin 2020;70:321-46.

22

Screening for Cervical Cancer*

Comparison of Screening Guidelines for Individuals as Average Risk for Cervical Cancer

Age	2024 USPSTF (draft recommendation)	2020 ACS and 2026 Update
< 21 years	Recommends against screening	No screening is recommended
21-24 years	Cytology alone every 3 years	No screening is recommended
25-29 years	Cytology alone every 3 years	- Primary HPV testing every 5 years with clinician-collected sample or every 3 years with self-collected sample - Cotesting* every 5 years or cytology alone every 3 years are acceptable options if primary HPV screening is limited or not available
30-65 years	- Primary HPV screening every 5 years with clinician- or patient-collected sample - Cotesting* every 5 years or cytology alone every 3 years is an alternative to primary HPV screening	- Primary HPV screening every 5 years with clinician-collected sample or every 3 years with self-collected sample - Cotesting* every 5 years or cytology alone every 3 years is acceptable if primary HPV testing is limited or not available
> 65 Years	Recommends against screening in those who have had adequate screening and are not otherwise at high risk for cervical cancer	Discontinue screening if negative primary HPV screening (preferred) or negative cotesting (acceptable) at 60 and 65 years of age, or three consecutive negative cytology tests performed at the recommended screening interval with last test at age 65 years (acceptable) - Self-collected primary HPV vaginal specimens should follow 3-year testing interval
After hysterectomy	Recommends against screening after hysterectomy with removal of the cervix and no history of CIN2+ or cervical cancer	Individuals without a cervix and without a history of CIN2+ in the past 25 years or any history of cervical cancer should not be screened.

**Recommendations apply to all women regardless of sexual activity status.*

23

US Health Resources and Services Administration January 2026

Approved Update Guidelines from the Women's Preventive Services Initiative

- *Designate high-risk HPV testing as the preferred screening modality for women at **average risk who are 30-65 years of age***

Result

- *Private Insurance Companies are **REQUIRED** to cover this preventive service **WITHOUT** cost-sharing by 1 January 2027*

[Update to the Women's Preventive Services Guidelines](#)

24

Self-Collected Vaginal Specimens for HPV Testing: Recommendations from the Enduring Consensus Cervical Cancer Screening and Management Guidelines Committee (2025)


Recommendation	
# 1	Clinician-collected cervical specimens are preferred, and self-collected vaginal specimens are acceptable for cervical cancer screening
# 2	When self-collected vaginal specimens are HPV-negative in the screening setting, repeat testing in 3 years is recommended
# 3	When self-collected vaginal specimens are positive for HPV 16 and/or 18, direct referral for colposcopy with concurrent cytology collection is recommended
# 4	When self-collected vaginal specimen HPV test results are: a) positive for HPV (untyped), b) negative for HPV 16/18 and positive for HPV HR12 (other); or c) negative for HPV 16/18 and positive for HPV 45, 33/58, 31, 52, 35/39/68, 51 or combinations thereof, obtaining a clinician collected cervical specimen for cytology or dual stain is recommended. Subsequent management of cytology or dual-stain results per management guidelines is recommended

Wentzensen N, Massad LS, Clarke MA, Garcia F, Smith R, Murphy J, Guido R, Reyes A, Phillips S, Berman N, Quinlan J, Lind E, Perkins RB; Enduring Consensus Cervical Cancer Screening and Management Guidelines Committee. Self-Collected Vaginal Specimens for HPV Testing: Recommendations From the Enduring Consensus Cervical Cancer Screening and Management Guidelines Committee. *J Low Genit Tract Dis.* 2025 Apr 1;29(2):144-152. doi: 10.1097/LGT.0000000000000885. Epub 2025 Feb 21. PMID: 39982254; PMCID: PMC11939108.

25

Harper DM, Bettcher CM, Young AP. It Is Time to Switch to Primary HPV Screening for Cervical Cancer. *Am Fam Physician.* 2024 Jan;109(1):8-9. PMID: 38227863.

Risk Management of Primary HPV Screening Results Using Genotyping Triage (Preferred)




HPV = human papillomavirus.

26

Harper DM, Bettcher CM, Young AP. It Is Time to Switch to Primary HPV Screening for Cervical Cancer. *Am Fam Physician*. 2024 Jan;109(1):8-9. PMID: 38227863.

Risk Management of Primary HPV Screening Results When Genotyping Is Unavailable



27

Important Considerations

No Screening

Women greater than 65 with a history of regular normal results


After hysterectomy for any condition
OTHER THAN Cervical cancer

Don't screen with Pap cytology if under 21 years of age and pap cytology or HPV testing if over 69 years of age

Individual Risk Factors: These are general guidelines. Individual risk factors (e.g., history of abnormal results, HPV infection, weakened immune system, DES exposure) may warrant more frequent screening.

Vaccination Status: Current guidelines recommend following age-based screening recommendations regardless of HPV vaccination

28



29

Primary HPV Testing *Bottom Line*

- Primary HPV screening has been thoroughly tested in 13 population-based randomized controlled trials over 15 years of follow-up.
- All trials found that primary HPV screening is as effective at detecting incident CIN 3+ as cotesting, with fewer harms.

Screening for Cervical Cancer

30

AVAILABLE HPV TESTS

31

HPV Tests FDA Approved for - ☒ HPV Primary Screening, ☒ Co-Testing, and ☒ Reflex Testing for ASC-US Results

Provides individual, specific results for **HPV 16, HPV 18, and HPV 45**, while grouping the remaining 11 genotypes.

Individual results of 6 high-risk HPV						Grouped results of 8 moderate risk HPV		
16	18	45	31	51	52	33/58	35/39/68	56/59/66
✓	✓	✓	✓	✓	✓	✓	✓	✓

12 pooled hrHPV

31	33	35	39	45	51
52	56	58	59	66	68

HPV16 HPV 18



Roche cobas HPV Assay

Abbot Alinity m High Risk HPV Assay

BD Onclarity HPV Assay

All three are available from the two most common Pap vials used by labs in the US -

- BD SurePath Liquid-based Pap Test
- Hologic ThinPrep Pap Test

32

What Can We Say?

Although primary HPV screening is as effective as cotesting at detecting cervical cancer, primary HPV screening decreases the number of lifetime screenings needed.

USPSTF has modeled different screening strategies, defining harms as the lifetime number of tests, colposcopies, false-positive results, cervical cancer cases, and cervical cancer deaths.

• Benefits were defined as life-years gained, disease detected, and cancer averted.

33

Lifetime Screening Outcomes for Three Cervical Cancer Screening Strategies per 1,000 Women

Strategy	Cytology tests	HPV tests	Total tests*	Colposcopies	Total tests/ CIN3+	Colposcopies/ CIN3+	Total workup/ CIN3+
Cyto-3y,21/ COTEST-5y,30	11,425	8,380	19,806	1,630	366.8	30.2	397
Cyto-3y,21/ HPV-5y(16/18),30	3,675	8,476	12,151	1,635	229.3	30.8	260
Cyto-3y,21/ HPV-5y(cyto),30	3,888	8,459	12,348	1,452	233.0	27.4	260

CIN 3+ = cervical intraepithelial neoplasia grade 3 (invasive squamous cell carcinoma, invasive adenocarcinoma, invasive adenosquamous carcinoma); cyto = cytology test; HPV = human papillomavirus; y = years of age.

*—Total number of cytology tests, HPV tests, and colposcopies performed to find CIN 3+.

Information from Kim JJ, Burger EA, Regan C, Sy S. Screening for cervical cancer in primary care: a decision analysis for the US Preventive Services Task Force. JAMA. 2018;320(7):706-714.

- Modeling the screening strategy of cervical cytology alone every three years for women 21 to 29 years of age, followed by cotesting every five years for women 30 to 65 years of age, led to the highest number of lifetime cytology tests per 1,000 women.
- For the individual, primary HPV screening provides equally accurate disease detection with fewer tests.

Harper DM, Bettcher CM, Young AP. It Is Time to Switch to Primary HPV Screening for Cervical Cancer. Am Fam Physician. 2024 Jan;109(1):8-9. PMID: 38227863.

34

Primary HPV Testing

• 2020's

- Primary HPV screening has been thoroughly tested in 13 population-based randomized controlled trials over 15 years of follow-up.
- All trials found that primary HPV screening is as effective at detecting incident CIN 3+ as cotesting, with fewer harms.

Screening for Cervical Cancer

35

Collection/Tests

e.g., Quest Diagnostics

Collection Is Exactly How
You Have ALWAYS Done It

Test Code 14264	HPV DNA (16, 18, Other High Risk), PCR, Cervical This test is used for the detection of high-risk Human Papillomavirus in clinician-collected cervical samples and can be used as part of an HPV primary screening algorithm. The test detects high-risk HPV types 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66, and 68, which are reported out as HPV 16, HPV 18, and other high-risk HPV types which includes, but does not differentiate, the remaining 12 high-risk types. Interim guidance...	⋮
Test Code 93296	HPV DNA (16,18, Other High Risk), PCR with Reflex to ThinPrep® Automated Pap	⋮
Test Code 14263	HPV DNA (16, 18, Other High Risk), PCR, Vaginal Self-Collected This test is used for the detection of high-risk Human Papillomavirus in self-collected vaginal samples and can be used as part of an HPV primary screening algorithm. The test detects high-risk HPV types 16, 18, 31, 33, 35, 39, 45, 51, 52, 56, 58, 59, 66, and 68, which are reported out as HPV 16, HPV 18, and other high-risk HPV types which includes, but does not differentiate, the remaining 12 high- risk types. Interim ...	⋮

36

Primary HPV Testing Effective Screening Modality for Cervical Cancer in Average-Risk Women

Risk Stratification Options for Cervical Cancer Screenings

Population	Recommendation
Low risk Hysterectomy including the cervix (exception cervical cancer; HGSIL) Cervix with comorbidities and decreased life expectancy	No testing
Average risk Patients (25 or) 30 to 65 years of age with a cervix	Primary HPV testing
High risk* Patients 30-65 years of age AND a high-risk condition (i.e., solid organ transplantation, chronic diseases marked by Janus kinase inhibitors or biologics, immunosuppressive drug treatment, HIV/AIDS postcoital bleeding, or under surveillance for prior HPV+ testing per ASCCP guidelines)	Cytology + HPV testing

* - High risk is NOT defined by smoking status, drug use, # of sex partners, or age.

Harper DM, Bettcher CM, Young AP. It Is Time to Switch to Primary HPV Screening for Cervical Cancer. Am Fam Physician. 2024 Jan;109(1):8-9. PMID: 38227863.

37

INTEGRATION OF DUAL-STAIN BIOMARKERS EXTENDED HPV GENOTYPING

38

Dual-Stain Biomarkers Robust Marker of CIN 3+

- Early 2024 – formal recognition of dual-stain cytology as an important tool for triaging HPV positive cervical cancer screening results.
- Guidelines include dual stain testing as an option in BOTH HPV primary screening and co-testing pathways.




*Technology detects the simultaneous presence of p16 and Ki-67 biomarkers in a single cell, which is a strong indicator of an HPV infection **progressing to precancer**.*

39

HPV Genotyping and p16/Ki-67 Test Significantly Improve Detection Rate of High-Grade Cervical Squamous Intraepithelial Lesion

Mechanism: Identifies the simultaneous expression of p16 (cell cycle arrest) and Ki-67 (cell proliferation) within a single cell—a **hallmark of HPV-mediated oncogenic transformation**.

Clinical Utility: For patients who are **hrHPV (+)** but have **Normal Cytology**, a positive Dual Stain can "up-triage" them to immediate colposcopy rather than waiting 12 months, as it is more specific for high-grade lesions.



- Brown cytoplasmic chromogen corresponds to p16 overload (white arrow)
- Nuclear compartment red chromogen is a proof of high Ki-67 protein accumulation and proliferative activity (black arrow)
- Red arrow depicts unaffected cell (case of HPV multi-infection), magnification 40×

DOI: 10.5114/aoms.2018.80697
Arch Med Sci 2020;16(1):87-93
With permission

Archives of
MEDICAL SCIENCE

40

Aim of Dual-Stain Integration

- Improve early diagnosis of HPV-positive cervical precancer and cancer
- Potentially reduce the number of women diagnosed with cervical cancer
- Require fewer colposcopies compared to cytology alone for HPV+ women
- Allow clinicians to better differentiate between HPV infections that may resolve on their own and those that require intervention

41

2024 ASCCP Enduring Risk-Based Management Guidelines

Aim to Integrate New Technologies and Approaches that Improve Cancer Prevention for High-Risk Individuals and Decrease Unnecessary Procedures in Lower-Risk Individuals

Extended Genotyping

Defined as the identification of individual types or groups of types beyond just HPV 16, 18 and 45. The new 2024 guidelines include HPV genotyping beyond HPV 16/18, to include management recommendations for HPV 31, 45, 33/58, 52, 35/39/68, 51 and 59/56/66.

Provides a greater and more refined level of risk stratification than pooled HPV-positive results to more precisely determine patient management.

HPV persistence (repeated detection of the same HPV type) is associated with a higher risk of cervical precancer compared to a single time point detection.


Unnecessary procedures can be safely decreased in lower-risk individuals who test positive for HPV 56/59/66

[file:///Users/davidweismiller/Downloads/Enduring%20Guidelines%20Extended%20Genotyping%20Summary%20032224%20\(2\).pdf](file:///Users/davidweismiller/Downloads/Enduring%20Guidelines%20Extended%20Genotyping%20Summary%20032224%20(2).pdf)

42

Primary HPV Testing

Refines Patient Risk Stratification and Advances the Way in Which We Screen for Cervical Cancer.



43

DEVELOP A FRAMEWORK FOR IMPLEMENTING PRIMARY HPV TESTING IN CLINICAL PRACTICE, INCLUDING PATIENT EDUCATION AND FOLLOW-UP STRATEGIES

44

ARS Question #2

Regarding Self-Collection of Vaginal Samples for HPV Testing in Cervical Cancer Screening, Which of the Following Statements Is True?

- A. Self-sampling has the potential to decrease the number of women who complete cervical cancer screening.
- B. Vaginal self-sampling has demonstrated slightly inferior sensitivity compared to clinician-collected cervical os sampling for HPV detection.
- C. Self-collection for HPV testing is currently only FDA approved for use both in a health care setting and at home.
- D. The vaginal self sampling approach for HPV is more effective at detecting HPV than a first void urine.

New Screening Landscape



"The cervical cancer screening landscape is rapidly changing. The use of vaginal samples, self-collected in a healthcare setting, marks a significant milestone in our joint efforts to improve accessibility and equity in healthcare."

Self-Collection

- Self-sampling is reasonable because HPV is transmitted skin-to-skin and must travel through the vagina before reaching the cervix.
- Two self-sampling approaches have been found to be effective.
 - Vaginal approach uses a device similar to a thin tampon that the patient inserts into the vagina, turns several times, places in a transport tube (some with liquid media), and returns.
 - Urine approach is similar to collection for sexually transmitted infection testing by gathering the desquamated cells coming from the cervix, vagina, and external genitalia in the first part of the urine stream.
- **Both techniques** are noninferior to clinician sampling at the os.

47



- Self-collection can improve cervical cancer screening access, especially in underserved populations.
 - In the US, Black, Hispanic and American Indian women have higher rates of cervical cancer than women of other racial groups; Black women having the highest rate of death.
- With vaginal self-collection as an option, women are more inclined to participate in such care, allowing health care providers an alternative option to identify a high-risk HPV infection in more convenient care settings.

CDC National Health Interview Survey (NHIS) 2023; Smith et al., J Clin Epidemiol. 2021; and CDC/NIH Behavioral Risk Factor Surveillance System on screening barriers.

48

Benefits to Self-Collection



- Self-sampling promotes health equity by allowing more women to obtain their own sample
- Potentially increasing the number of women who complete screening.



- Early cost-effectiveness analyses suggest overall cost savings
- More importantly, that the savings from not performing cotesting could cover current out-of-pocket costs for colposcopy.



- The Affordable Care Act mandates coverage only for screening tests.
- U.S. Congress could amend it to cover all screening and evaluation costs for cervical cancer, payors could easily shift the costs from unnecessary cytology screening to cover the out-of-pocket costs for colposcopy, as they have done for colorectal cancer screening.

49

The STEP Randomized Clinical Trial (2026 Analysis)

- New economic and longitudinal data from the **STEP trial** (Self-Testing options in the Era for Primary HPV screening) was published in early 2026.
- **Key Finding**
 - Directly mailing HPV kits to overdue individuals **increased screening rates by over 14%**.
 - Analysis confirmed this approach is not only cost-effective but **cost-saving** compared to traditional office-based Pap cytology.

*Economic Analysis of Mailed Primary HPV Self-Testing Kits in a Large Integrated Health System.
JAMA Network Open (March 2026).*

50

Primary HPV Screening Tests Approved by the FDA

Test (year approved)	Genotyping	Approved for Self Collection
Roche cobas HPV test (2014)	HPV 16 and 18 Detects 12 other high-risk HPV types: 31/33/35/39/45/51/52/56/58/59/66/68 (pooled results)	Yes, using Evalyn brush, Copan 522C.80 swab (FLOQSwab), or Teal Wand
BD Onclarity HPV Assay (2018)	HPV 16, 18, and 45 Detects 11 other high-risk HPV types: 31, 51, 52, 33/58, 35/39/68, and 56/59/66 (pooled results)	Yes, using Copan 522C.80 swab
Abbott Alinity m High Risk HPV Assay (2023)	HPV 16, 18, and 45 genotyping not provided but can be added as additional testing; 11 others can also be added: 31/33/52/58 and 35/39/51/56/59/66/68	Yes, using simpli-COLLECT HPV Collection Kit or Evalyn brush

51

Primary HPV Screening Tests Approved by the FDA

Test (year approved)
Roche cobas HPV test (2014) Evalyn brush, Copan 522C.80 swab (FLOQSwab), or Teal Wand
BD Onclarity HPV Assay (2018) Copan 522C.80 swab
Abbott Alinity m High Risk HPV Assay (2023) simpli-COLLECT HPV Collection Kit or Evalyn brush



52

At-Home Cervical Cancer Screening Kit & Virtual Visit

Same HPV test and accuracy as screening in-person, but now you can collect your own sample, privately from home.

To get early access, enter your email below.

\$99 with insurance
\$249 without insurance (reduced from \$499)

Enter email address*

Not due yet? [Schedule a reminder](#)

Teal Wand

*Clinical Sensitivity – 96% (Equal)
to That of Clinician Collection*

F.D.A. Approves First At-Home Alternative to the Pap Smear

The tool will allow women to screen for HPV, which causes almost all cases of cervical cancer, without visiting a doctor.

Wall Street Journal May 9, 2025

Simple 3 Step Process

Get the same accurate screening you'd receive at a clinic, but from the comfort of home - with support available every step of the way.

Step 1	Step 2	Step 3
Order & Attend Consult Request a kit and attend a short visit* with a Teal provider to review your screening history and learn about screening at home.	Receive Kit & Collect Receive your kit, comfortably collect your sample, and mail it to the lab.	Results & Doctor Support View results in your Teal portal. If follow-up care is needed, you can connect virtually with a Teal provider to discuss next steps and coordinate your referral care.
⌚ 1-3 days	⌚ 3-5 days	⌚ 4-5 days

*The Teal Wand requires a prescription, so a short virtual visit with a Teal provider is needed before we can mail your kit.

53

ACS – 2025 Teal Wand

American Cancer Society
Includes Teal's At-Home Self-Collection for Primary HPV Testing in New Cervical Cancer Screening Guidelines

December 4, 2025



(2026), Screening for cervical cancer. CA Cancer J Clin, 76: e70049. <https://doi.org/10.3322/caac.70049>

54

David Glenn Weismiller, MD
Goodbye Pap, Hello Primary HPV Testing

Teal Health



Patients will label a vial with the sample in it, and then mail it out for processing.



After a sample is collected, the swab is detached from the wand and returned in the vial.

- Allow patients to order the test online, speak with a telehealth provider, collect a sample and then mail it to be tested for HPV.
- Company is working with insurers to secure coverage and with donors to try to subsidize the cost for people without insurance.
- If the test is positive, the patient will be referred to an in-person provider for a cytologic sampling or colposcopy.
 - Teal plans to mail an initial round of tests to customers in California in June before expanding to other states.

55

HPV DNA (16, 18, Other High Risk), PCR, Vaginal Self-Collected

Test Code

14263  

CPT Code(s)*

87626

Test Details

Methodology

Polymerase Chain Reaction (PCR)

Reference Range(s)

HPV 16	Not Detected
HPV 18	Not Detected
HPV Other	Not Detected

Alternative Name(s)

Primary Screening, HPV alone, HPV Vaginal Collection, HPV self-collect

Preferred Specimen(s)

1 ThinPrep® vial sample collected with COPAN FLOQSwab 552C.RM

Alternative Specimen(s)

1 ThinPrep vial sample collected with Evalyn Brush

Minimum Volume

1.5 mL

Collection Instructions

Patient should follow instructions provided with the self-collection device.

The collection device should be immediately transferred to ThinPrep following provided instructions.

Transport Container

ThinPrep® vial

Transport Temperature

Room temperature

e.g., **Quest Diagnostics**

56

Let's Get Real...



Some real head butting over this...


57

Let's Get Real...

- Cervical cancers missed with HPV-Alone*

1 in 5

* No one ever said do this alone. A (+) screening result may lead to further evaluation with cytology and /or colposcopy



Among **adequately screened women**,


HPV self-collect was the **least preferred** method of screening.

Providers Beliefs

- Fewer providers **believed** in the effectiveness of pHPV to reduce cervical cancer mortality and were less likely to recommend pHPV in the correct age group and screening interval compared with cytology-based screening modalities

Kruse G et. al. JNCI Cancer Spectrum. 2023;7(1)

58



The image shows the ASCCP website header with the logo and navigation menu. The main heading is "ASCCP 2019 Risk-Based Management Consensus Guidelines". Below this, a paragraph describes the guidelines for abnormal screening results. A button labeled "Download Guidelines" is present. A large blue banner with the text "LIVING GUIDELINES" is followed by a paragraph explaining the shift in management for primary HPV testing from test results to individualized risk-based triage.

ASCCP Improving lives through the prevention and treatment of anogenital & HPV-related diseases

Log In Create an Account

Search... Donate Mobile App

Guidelines Practice Management Education & Events Journal Membership News Partners About

ASCCP 2019 Risk-Based Management Consensus Guidelines

Risk-based Management Guidelines for abnormal screening results, including post-colposcopy or post-treatment scenarios. Enduring Guidelines apply this approach to new technologies that emerge and novel clinical scenarios that were not previously addressed.


ASCCP 2019 Risk-Based Management Consensus Guidelines

Download Guidelines

LIVING GUIDELINES

*The 2025–2026 management landscape for **Primary HPV testing** has shifted from "test results" to **individualized risk-based triage**. The core principle is "Equal Risk, Equal Management," where a patient's prior history now carries as much weight as their current HPV status.*

59



The image is a banner for the ASCCP Risk-Based Management Consensus Guidelines app. It features a laptop and a smartphone displaying the app interface. The text "ASCCP Risk-Based Management Consensus Guidelines" is prominently displayed. Below it, the text "Download the App" is followed by buttons for "Available on the App Store", "GET IT ON Google Play", and "Web Application". A red arrow points to the "Web Application" button. At the bottom, there is a link for "Need Help? Quick Start User Guide".

ASCCP Risk-Based Management Consensus Guidelines

Download the App

Available on the App Store GET IT ON Google Play Web Application

Need Help? Quick Start User Guide

60

ASCCP Living Guidelines (Risk-Based Management)

We no longer treat the result; we treat the risk -

- **What are "Living" Guidelines?** The ASCCP moved to a web-based/app-based model because the data is updated in real-time. It uses the "Equal Management for Equal Risk" principle.
- **The 4% Threshold:** If a patient's 5-year risk of CIN 3+ is < 4%, we can usually follow up in 1, 3, or 5 years. If the risk is $\geq 4\%$, a Colposcopy is recommended.
- **Why it Matters:** Prevents over-treatment in low-risk patients (like young women with transient HPV) while catching high-risk patients who might have had a "negative" Pap but carry HPV 16 or 18.
- **History is important** – it is ideal to have 10 years, 5 years works.


61

Application

The screenshot displays the 'Management' tab of the ASCCP Living Guidelines application. The interface includes a sidebar labeled 'Application' and a main content area with three sections: 'AGE', 'CLINICAL SITUATION', and 'SPECIAL SITUATIONS'. Each section has a red arrow pointing to it from the left. The 'AGE' section has three buttons: 'Under 25', '25 to 29', and '30+'. The 'CLINICAL SITUATION' section has four buttons: 'Routine screening', 'Evaluation of a colposcopic biopsy', 'Management of results during post colposcopy surveillance (within past 7 years)', and 'Follow-up after treatment'. The 'SPECIAL SITUATIONS' section has four buttons: 'Unsatisfactory test results', 'Post hysterectomy', 'Symptomatic patients', and 'Immunosuppressed patients'.

62

Web Application



The image shows two screenshots related to the ASCCP Management Guidelines Web Application. On the left is the registration form with fields for First Name, Last Name, and Email, along with a checkbox for newsletters and a 'Next >' button. On the right is a confirmation email from info@asccp.org via amazonSES.com, dated 3:17 AM. The email contains a 'Welcome' message and a 'Get started' button, which is highlighted with a red arrow.

63

The “Treat” Threshold

- When do we skip the biopsy and go straight to the LEEP?
 - **Expedited Treatment (See-and-Treat):** If the immediate risk of CIN 3+ is >25% (e.g., HPV 16 positive with HSIL cytology), the guidelines now allow for moving directly to treatment (LEEP) without a biopsy first, especially in patients over 25 who have completed childbearing.
 - **The Goal:** Reduce the "loss to follow-up" an

64

Summary of Management for Positive HPV Screening Test Results with Self-Collected and Clinician-collected Specimens for Settings Using Cytology (Either as a Cotest or Triage Test)

ASCCP

65

HPV test result	Management of clinician- vs. self-collected collected specimens	Current HPV test result	Current cytology result	Past history	Management
HPV 16/18	Clinician-collected: Laboratory performs cotest or reflex cytology. Self-collected: Colposcopy recommended. Collect cytology at colposcopy.	16	HSIL	Noncontributory	Treatment preferred; colposcopy acceptable
		16	ASC-H	Noncontributory	Treatment or colposcopy
		16	NILM, ASC-US, LSIL, AGC, or no cytology	Noncontributory	Colposcopy ¹
		18	HSIL	Noncontributory	Treatment or colposcopy
		18	NILM, ASC-US, LSIL, ASC-H, AGC, or no cytology	Noncontributory	Colposcopy ^{1,2}
HPV 45, 33/58, 31, 52, 35/39/68, 51 Untyped or "other" types when 16 and 18 are not present	Clinician-collected: Laboratory performs cotest or reflex cytology. Self-collected: Patient returns for collection of cytology unless current test is 2 nd consecutive HPV+ in which case colposcopy recommended.	45, 33/58, 31, 52, 35/39/68, 51 or untyped/other	HSIL, ASC-H, AGC	Noncontributory	Colposcopy ^{1,2}
		45, 33/58, 31, 52, 35/39/68, 51	ASC-US, LSIL	Noncontributory	Colposcopy
		Other/untyped	ASC-US, LSIL	Documented HPV negative screen in past 5 years or colposcopy <CIN2 within past 1 year	Repeat HPV test in 1 year
		Other/untyped	ASC-US, LSIL	Any history other than above	Colposcopy
		45, 33/58, 31, 52, 35/39/68, 51 or untyped/other	NILM	Normal ³ or colposcopy <CIN2 within past 1 year	Repeat HPV test in 1 year
		45, 33/58, 31, 52, 35/39/68, 51 or untyped/other	Not available	HPV+ without colposcopy (i.e. current test is 2 nd consecutive HPV+)	Colposcopy
HPV 59/56/66	Clinician-collected: No additional immediate testing needed. Laboratory may run cytology if cotesting is performed. ⁴ Self-collected: No additional immediate testing needed	59/56/66	ASC-H, AGC, or HSIL	Noncontributory	Colposcopy ^{1,2}
		59/56/66	No cytology or NILM, ASC-US, LSIL	Normal or colposcopy <CIN2 within past 1 year	Repeat HPV test in 1 year
		59/56/66	Not available	HPV+ without colposcopy (i.e., current test is 2 nd consecutive HPV+)	Colposcopy

Wentzensen N, Massad LS, Clarke MA, Garcia F, Smith R, Murphy J, Guido R, Reyes A, Phillips S, Berman N, Quinlan J, Lind E, Perkins RB; Enduring Consensus Cervical Cancer Screening and Management Guidelines Committee. Self-Collected Vaginal Specimens for HPV Testing: Recommendations From the Enduring Consensus Cervical Cancer Screening and Management Guidelines Committee. J Low Genit Tract Dis. 2025 Apr 1;29(2):144-152. doi: 10.1097/LGT.0000000000000885. Epub 2025 Feb 21. PMID: 39982254; PMCID: PMC11939108

66

Summary of Management for Positive HPV Screening Results with Self-collected and Clinician-collected Specimens for Settings Using Reflex Testing with p16ink4a/Ki-67 Dual Stain

ASCCP

67

HPV test result	Management of clinician- vs. self-collected collected specimens	Current HPV result	Current dual stain result	Past history	Management
HPV 16/18	Clinician- and self-collected: same management	16 and/or 18	Noncontributory	Noncontributory	Colposcopy with collection of cytology if available
HPV 45, 33/58, 31, 52, 35/39/68, 51 or untyped	Clinician-collected: Laboratory performs reflex dual stain. Self-collected: Patient returns for collection of dual stain.	45, 33/58, 31, 52, 35/39/68, 51 or untyped	Dual stain negative ¹	Normal ² or colposcopy <CIN2 within past 1 year	Repeat HPV test in 1 year
		45, 33/58, 31, 52, 35/39/68, 51 or untyped	Dual stain positive ³	Noncontributory	Colposcopy
		45, 33/58, 31, 52, 35/39/68, 51 or untyped	Noncontributory	HPV+ without colposcopy (i.e., current test is 2 nd consecutive HPV+)	Colposcopy
HPV 59/56/66	Clinician- and self-collected: same management	59/56/66	Noncontributory	Normal ² or colposcopy <CIN2 within past 1 year	Repeat HPV test in 1 year ¹
		59/56/66	Noncontributory	HPV+ without colposcopy (i.e., current test is 2 nd consecutive HPV+)	Colposcopy

Wentzensen N, Massad LS, Clarke MA, Garcia F, Smith R, Murphy J, Guido R, Reyes A, Phillips S, Berman N, Quinlan J, Lind E, Perkins RB; Enduring Consensus Cervical Cancer Screening and Management Guidelines Committee. Self-Collected Vaginal Specimens for HPV Testing: Recommendations From the Enduring Consensus Cervical Cancer Screening and Management Guidelines Committee. *J Low Genit Tract Dis*. 2025 Apr 1;29(2):144-152. doi: 10.1097/LGT.0000000000000885. Epub 2025 Feb 21. PMID: 39982254; PMCID: PMC11939108.

68

David Glenn Weismiller, MD
Goodbye Pap, Hello Primary HPV Testing

Key 2026 Action Thresholds

Management decisions are now mathematically guided by "Immediate Risk" of CIN3+:

Risk	Management
≥ 60%	Expedited treatment (LEEP) is preferred.
25-59%	Expedited treatment or colposcopy is acceptable.
4.1-24%	Colposcopy is required
0.15-4%	1-year follow-up.
< 0.15%	Return to 5-year screening.

*These thresholds are calculated dynamically via the **ASCCP Management Guidelines App**, which was updated in January 2026 to include longitudinal data from over 1 million patients in the Kaiser Permanente-Northern California cohort.*

69

The Script for the Skeptical Patient

"I know it feels strange to skip the Pap smear, but we've actually upgraded our technology. We are now testing for the HPV virus itself rather than just looking for the cell changes it causes. Think of it like a security system: the old Pap smear looked for 'signs of a break-in' (cell changes), while the new test checks to see if 'the intruder is even at the door' (the virus). Because this test is so much more accurate at predicting your risk over the next several years, it is actually safer to wait longer between tests. We are still doing your pelvic exam and health check today—we're just being smarter about how we screen for cancer."

70

RETURN TO OUR CASES

71

The Clinical Conundrum Question 3

Patient: Sarah, a 31-year-old G2P2

History: No prior abnormalities; last screen 3 years ago was a "Normal" Pap (no HPV).

Today's Result: Primary HPV is **Positive** (Non-16/18). Reflex Cytology is **Negative** (NILM).

The Patient's Question: *"Wait, you found the virus that causes cancer, but you say my cells are normal? Am I sick or not? And what do we do now?"*



CONTINUING EDUCATION COMPANY

72

...Response?

Primary HPV or co-testing setting patients positive for HPV 45, 33/58, 31,52, 35/39/68, 51 or combination thereof, but negative for HPV16 and HPV 18.

*recommend Clinician Collected Primary HPV Screening in One Year

Triage with cytology or dual stain (DS) is recommended.

- If DS (-) or NILM cytology, repeat testing in 1 year
- If DS (+) or cytology $\geq$ ASC-US, colposcopy recommended.

73

ARS Question #3

Based on the 2026 ASCCP "Living" Guidelines, What Is the Most Appropriate Next Step for Sarah?

- A. Immediate Colposcopy
- B. Repeat co-testing (HPV + Pap) in 1 year
- C. Repeat Primary HPV testing in 1 year
- D. Pap Cytology with reflex HPV in 1 year

74

Patient Data

Screening Management Publications Definitions

CURRENT HPV

HPV 16+

HPV 18+

HPV+ (untyped)

HPV+ (31/33/35/39/45/51/52/58/68)

HPV+ (16/18 negative, other high risk positive)

HPV+ (56/58/66)

HPV-

None

CURRENT RESULTS

A Positive HPV Result needs further testing for triage. What other results does this patient have, if any?

☒ Normal

☐ ASC-US

☐ LSIL

☐ ASC-H

☐ HSIL

☐ AGC

☐ Dual Stain Negative

☐ Dual Stain Positive

PRIOR HISTORY

Does the patient have abnormal results on the most recent prior screening? Leave blank if history is unknown or if last screen was more than approximately 5 years ago.

- ☐ HPV Positive
- ☐ HPV Negative
- ☒ Normal
- ☐ ASC-US
- ☐ LSIL
- ☐ ASC-H
- ☐ HSIL
- ☐ AGC

Screening Management Publications Definitions

2 AGE

Under 25 25 to 29 30+

3 CLINICAL SITUATION

Routine screening

Evaluation of a colposcopic biopsy

Management of results during post colposcopy surveillance (within past 7 years)

Follow-up after treatment

4 SPECIAL SITUATIONS

Unsatisfactory test results

Post hysterectomy

Symptomatic patients

Immunosuppressed patients

<https://www.asccp.org/mobile-app/>

75

Recommendation

CONFIRMATION

Age: 30+
Clinical Situation: Routine screening

CURRENT RESULTS

Cotest with a positive HPV test and normal cytology
Extended genotyped (other high-risk HPV+)

PRIOR RESULTS

Cytology-only test with a normal result


SCREENING RECOMMENDATION

HPV Positive Normal Cytology

1 year follow up¹

HPV-based screening at follow-up visit; clinician collected preferred.^{2,3}

Risk



76

The Answer and Rationale

- **Correct Answer:**

- **The 4% Rule:** Sarah's immediate risk of CIN 3+ is well below the 4% threshold required for a colposcopy.
- **The "Why":** Most HPV infections in healthy 30-somethings are transient. By waiting 12 months, we allow for "viral clearance."
- **The Nuance:** If we did a colposcopy today, we would likely find nothing or a low-grade lesion (LSIL/CIN 1) that would resolve on its own. We would be "over-diagnosing" and causing unnecessary cervical trauma.
- **The Next Step:** If she is still HPV positive in one year, her risk rises above the 4% threshold, and then we perform the colposcopy.

Sarah is the most common patient in your 2026 inbox. If you send every Sarah to colposcopy, you'll have a backlog of procedures and a lot of anxious patients. If you use the ASCCP app, you give her a evidence-based 'green light' to wait for her immune system to do its job

77

The Clinical Conundrum Question 4

Patient: Maria, a 31-year-old G2P2

History: No prior abnormalities; last screen 3 years ago was a "Normal" Pap (no HPV).

Today's Result: Primary HPV is **Positive for HPV 16**

The Patient's Question: "Wait, you found the virus that causes cancer? Am I sick or not? And what do we do now?"



CONTINUING EDUCATION COMPANY

78

ARS Question #4

Based on the 2026 ASCCP "Living" Guidelines, What Is the Most Appropriate Next Step for Maria?

- A. Immediate Colposcopy
- B. Repeat co-testing (HPV + Pap) in 1 year
- C. Repeat Primary HPV testing in 1 year
- D. Expedited treatment (LEEP)



CONTINUING EDUCATION COMPANY

79

Data Entry

CURRENT HPV

HPV 16+

HPV 18+

HPV+ (untyped)

HPV+ (31/33/35/39/45/51/52/58/68)

HPV+ (16/18 negative, other high risk positive)

HPV+ (56/59/66)

HPV-

None

CURRENT RESULTS

A Positive HPV Result needs further testing for triage. What other results does this patient have, if any?

- ☐ Normal
- ☐ ASC-US
- ☐ LSIL
- ☐ ASC-H
- ☐ HSIL
- ☐ AGC
- ☐ Dual Stain Negative
- ☐ Dual Stain Positive

CONFIRMATION

Age: 30+

Clinical Situation: Routine screening

CURRENT RESULTS

HPV16 positive test without cytology or dual stain results

80

SCREENING RECOMMENDATION

HPV 16+ with No Cytology

Colposcopy with cytology^{1,2}

Cytology is recommended with Colposcopy if patient doesn't have cytology results.¹

When self-collected vaginal specimens are positive for HPV 16, direct referral for colposcopy with concurrent cytology collection is recommended.²

81

The Answer and Rationale

- **Correct Answer:**

- **The 4% Rule:** Maria's immediate risk of CIN 3+ is well above the 4% threshold required for a colposcopy.
- **The "Why":** the presence of **HPV 16** (or 18) carries an immediate risk of high-grade disease that is **higher than 4%**.

The genotype is our 'urgency' marker. Sarah and Maria have the exact same Pap result, but their clinical path is totally different because of the DNA of the virus they are carrying. That is the heart of 2026 precision screening.

The "Living" Guideline algorithm sees Maria's history as a sign of **persistent** HPV. Persistence is the true driver of cancer

82

The Clinical Conundrum 3

Patient: Sheila, a 57-year-old G3P3

- Establishing Care
- **History**
 - LNMP – 4 years ago
 - (+) VMS
 - Trial of Estrogen – no benefit
 - Recently had bioidentical Testosterone pellets implanted

- 8 years ago she underwent a hysterectomy for “precancer.”
- She reports she has never had an abnormal pap and she has been getting “yearly evaluations for as long as she can remember.”

The Patient’s Question – “*Can I move to primary HPV testing instead of Pap?*”

83

Summary Checklist for the Modern Clinician

We’ve Moved from 'What Does This Pap Mean?' to 'What Is This Patient's 5-Year Risk?' that Is a Massive Win for Patient Safety and Reducing Over-Treatment."

	Parameter
✓	Switch to Primary HPV: Start at age 25 if your lab supports it.
✓	Download the App: Use the ASCCP Management Guidelines app—don't try to memorize the tables.
✓	History is King: You cannot manage a current abnormal result without knowing the last 10 years of the patient's screening history.
✓	Genotyping Matters: 16 and 18 are the "bad actors." They always require closer scrutiny than other high-risk types.

84


“Cervical cancer screening recommendations have changed over time, driven by improved understanding of the disease's natural progression, the recognized role of high-risk HPV infection, and advancements in screening technologies”

SUMMARY

- Primary HPV screening benefits patients by limiting unnecessary testing, benefits physicians by providing a simpler strategy for management, and benefits society by making cervical cancer screening more accessible.
- Now is the right time for clinicians to shift our cervical cancer screening strategy from cotesting to primary HPV screening for average-risk, asymptomatic women.

85

Primary HPV Screening



86

The Challenge: *Changes in Practice*



Make primary HPV the default

Update EHR order sets; CPL, Quest, and LabCorp run an FDA-approved primary HPV assay. **We need to order one swab.**



Offer self-collection

For overdue patients and those who decline a speculum exam, in clinic or by mail



Close the loop on every positive

Track results in a registry; navigate patients to triage or colposcopy on time



Manage by risk

Use the ASCCP app; ask the lab for reflex extended genotyping and dual stain



Vaccinate at every visit

Start at 9–12, catch up through 26 (max 45), and track up-to-date status

Measure it: Audit primary HPV use, self-collection offers and on-time follow-up at baseline and 6 months

Practice Recommendations

Recommendation	SOR
Join the movement, increase access to cervical cancer screening. Use Primary HPV Testing. Contact your reference lab for collection specifics and ordering.	-
All trials found that primary HPV screening is as effective at detecting incident CIN 3+ as cotesting, with fewer harms.	A
Screening recommendations: (USPSTF (Draft) – 2024) (ACS- 2025)	
Ages 21-24: Cytology (Pap test) every 3 years	A
Ages 25-65: Primary HPV Testing every 5 years (clinician-collected) or 3 years (self-collected)	
Two Sampling approaches have been found to be effective for Primary HPV Testing self collection – Vaginal Approach and Urine (first catch) approach. Home self-collection is also now available.	B
Dual stain biomarkers integration allows clinicians to better differentiate between HPV infections that may resolve on their own and those that require intervention.	B
Extended genotyping provides a greater and more refined level of risk stratification (persistence) than pooled HPV-positive results to more precisely determine patient management.	B

References

1. Marcus, Jenna Z. MD¹; Cason, Patty RN, MS, FNP-BC²; Downs, Levi S. Jr. MD, MS³; Einstein, Mark H. MD, MS¹; Flowers, Lisa MD⁴. The ASCCP Cervical Cancer Screening Task Force Endorsement and Opinion on the American Cancer Society Updated Cervical Cancer Screening Guidelines. *Journal of Lower Genital Tract Disease* 25(3):p 187-191, July 2021. | DOI: 10.1097/LGT.0000000000000614
2. Perkins RB, Guido RS, Castle PE, Chelmow D, Einstein MH, Garcia F, Huh WK, Kim JJ, Moscicki AB, Nayar R, Saraiya M, Sawaya GF, Wentzensen N, Schiffman M; 2019 ASCCP Risk-Based Management Consensus Guidelines Committee. 2019 ASCCP Risk-Based Management Consensus Guidelines: Updates Through 2023. *J Low Genit Tract Dis.* 2024 Jan 1;28(1):3-6. doi: 10.1097/LGT.0000000000000788. PMID: 38117563; PMCID: PMC10755815.
3. Schiffman, Mark MD¹; Wentzensen, Nicolas MD¹; Perkins, Rebecca B. MD²; Guido, Richard S. MD³. An Introduction to the 2019 ASCCP Risk-Based Management Consensus Guidelines. *Journal of Lower Genital Tract Disease* 24(2):p 87-89, April 2020. | DOI: 10.1097/LGT.0000000000000531
4. Harper DM, Bettcher CM, Young AP. It Is Time to Switch to Primary HPV Screening for Cervical Cancer. *Am Fam Physician.* 2024 Jan;109(1):8-9. PMID: 38227863.
5. Self-collected vaginal specimens for human papillomavirus testing and guidance on screening exit: An update to the American Cancer Society cervical cancer screening guideline. *CA: A Cancer Journal for Clinicians* (2025).